

Essential



Zillarator

For Feedlot Cattle fed in confinement for slaughter

For increased rate of weight gain, improved feed efficiency and increased carcass leanness in cattle fed in confinement for slaughter during the last 20-40 days on feed

ACTIVE DRUG INGREDIENT

Zilpaterol Hydrochloride.....320 g/ton

GUARANTEED ANALYSIS

Crude Protein, min..... 5.00%

Crude Fat, min..... 2.50%

Crude Fiber, max.....30.00%

Calcium, min.....11.25% max..... 13.75%

FEEDING DIRECTIONS

Mix one (1) bucket (10 pounds) to 460 of complete cattle feed, to provide 6.8 grams of Zilpaterol hydrochloride per ton, or add ½ pound (1 rounded cup) to 23 pounds of grain. Cattle should consume 60-90 mg per head per day. Each pound contains 160 mg of Zilpaterol Hydrochloride.

Zillarator is fed the last 20-40 days prior to slaughter to cattle in confinement for increased weight gain, improved feed efficiency, and increased carcass leanness.

INGREDIENTS

Processed Grain Byproducts, Roughage products, calcium carbonate, Forage products, molasses products, animal fat, mineral oil, and Propionic Acid (as a preservative)

CAUTION: Do not allow horses or other equines access to feed containing Zilpaterol Hydrochloride. Not for use in animals intended for breeding. Do not use in veal calves.

Limitations for Use: Feed continuously as the sole ration during the last 20-40 days on feed. Withdrawal 3 days prior to slaughter

You May Notice: Animals receiving Zilpaterol Hydrochloride may exhibit increased respiratory rate as well as elevated levels of creatine phosphokinase (CPK) and creatinine.

Manufactured by: Nutrition Services, Inc.
501 Division Ave - York, NE 68467
(800) 697-1998

www.EssentialShowFeeds.com



Net Weight: 10 lbs. (4.53 kg)